

Rapidity Dependence of J/ψ Production at RHIC and LHC

Yunpeng Liu¹, Zhen Qu¹, Nu Xu², and Pengfei Zhuang¹

¹Physics Department, Tsinghua University, Beijing 100084, China

²Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

(Dated: July 16, 2009)

The motion of charmonium in heavy ion collisions is described by a three dimensional transport equation with initial production and continuous regeneration in hot medium. The observation of apparently stronger J/ψ suppression at forward rapidity compared to that at midrapidity, so called J/ψ puzzle at RHIC, can well be explained by the competition between the two production mechanisms. At LHC, however, the rapidity dependence of the J/ψ production is dominated by the regeneration process.

PACS numbers: 25.75.-q, 12.38.Mh, 24.85.+p

From lattice quantum chromodynamics (QCD) calculations, a new state of matter, the so-called quark-gluon plasma (QGP), has been predicted to exist at high temperature and/or high baryon density. The J/ψ production has long been considered as a probe of the QGP formation in relativistic heavy ion collisions [1]. Different from the J/ψ suppression observed at the CERN Super Proton Synchrotron (SPS) where almost all the charmonia are primordially produced through hard nucleon-nucleon collisions (NN) and then suffer from nuclear absorption [2, 3] in the initial stage and anomalous suppression in the hot and dense medium [4, 5, 6, 7, 8, 9] in the later stage, there are a remarkable number of charm quarks in the QGP produced in high energy nuclear collisions at the BNL Relativistic Heavy Ion Collider (RHIC) and the CERN Large Hadron Collider (LHC), and the regeneration, namely the continuous recombination of those uncorrelated charm quarks offers another source for J/ψ production [10, 11, 12, 13, 14]. Obviously, the regeneration will enhance the J/ψ yield and alter its momentum spectrum. At midrapidity, the initial production and regeneration are almost equally important and the competition between the two production mechanisms controls [15, 16] the J/ψ nuclear modification factor R_{AA} and averaged transverse momentum square $\langle p_t^2 \rangle$ observed at RHIC [17, 18].

Recently, the rapidity dependence of J/ψ production in Au+Au collisions is measured at RHIC [17, 19] and discussed in models [20, 21, 22]. As one can see, as a function of the number of participant nucleons N_p , the observed $R_{AA}(N_p)$ shows that the suppression is significantly stronger at forward rapidity ($|y| \in [1.2, 2.2]$) than that at midrapidity ($|y| < 0.35$). The enhanced suppression is especially clear in central collisions, see Figs. 4 and 5 of Ref. [17]. For $N_p > 150$, the ratio of $R_{AA}^{\text{forward}}/R_{AA}^{\text{mid}}$ is about 0.6. In addition, we observe that the value of $\langle p_t^2 \rangle$ is lower at midrapidity than that at forward rapidity, see Table II of Ref. [17]. The above observation is difficult to be explained by models with only initial charmonium production [17]. Since the medium temperature at midrapidity is higher than or at least the same as that at forward rapidity, the suppression at forward rapidity is predicted in these models to be less than or at most

the same with that at midrapidity.

In this Letter, we investigate the rapidity dependence of J/ψ production by constructing a three dimensional transport equation for the charmonium motion in the hot medium, with both initial production and regeneration. When the regeneration mechanism is included, the above J/ψ puzzle is possible to be explained [21]. Since the charm quarks are mainly distributed in the central region, their recombination into J/ψ s at midrapidity is more important than that at forward rapidity. While the medium absorption at midrapidity is in principle larger than that at forward rapidity, the competition between the regeneration and absorption may lead to an enhancement in R_{AA} at midrapidity. We note that the effect in the rapidity dependence of R_{AA} should also be elaborated with the rapidity dependence of $\langle p_t^2 \rangle$. Considering the fact that the measured forward rapidity is still located at the plateau of light hadron rapidity distribution [23], the temperatures of the medium and in turn the J/ψ absorption in the mid and forward rapidity are approximately the same. Therefore, the rapidity dependence of the regeneration becomes the key factor to control the experimental observation $R_{AA}^{\text{forward}}/R_{AA}^{\text{mid}} < 1$ at any N_p and $\langle p_t^2 \rangle^{\text{forward}}/\langle p_t^2 \rangle^{\text{mid}} > 1$ at large N_p at RHIC.

Since the experimental results can not separate primordial J/ψ and J/ψ from χ_c and ψ' decay, we consider a transport equation for a charmonium distribution function $f_\Psi(p, x)$ ($\Psi = J/\psi, \psi', \chi_c$). Using Lorentz covariant time $\tau = \sqrt{t^2 - z^2}$, space rapidity $\eta = 1/2 \ln[(t+z)/(t-z)]$, transverse energy $E_t^\Psi = \sqrt{E_\Psi^2 - p_z^2}$ with $E_\Psi = \sqrt{m_\Psi^2 + \mathbf{p}^2}$ and momentum rapidity $y_\Psi = 1/2 \ln[(E_\Psi + p_z)/(E_\Psi - p_z)]$, the three dimensional transport equation can be expressed as

$$\left[\cosh(y_\Psi - \eta) \frac{\partial}{\partial \tau} + \frac{\sinh(y_\Psi - \eta)}{\tau} \frac{\partial}{\partial \eta} + \mathbf{v}_t^\Psi \cdot \nabla_t \right] f_\Psi = -\alpha_\Psi f_\Psi + \beta_\Psi, \quad (1)$$

where $\mathbf{v}_t^\Psi = \mathbf{p}_t/E_t^\Psi$ is the transverse velocity which leads to the leakage effect and is proven to be important for those high velocity charmonia at SPS [9], and the suppression and regeneration in hot medium are described by the loss term α_Ψ and gain term β_Ψ . Considering

the gluon dissociation process $\Psi + g \rightarrow c + \bar{c}$ and charmonium regeneration process $c + \bar{c} \rightarrow \Psi + g$ in QGP, $\alpha(\mathbf{p}_t, y_\Psi, \mathbf{x}_t, \eta, \tau)$ and $\beta(\mathbf{p}_t, y_\Psi, \mathbf{x}_t, \eta, \tau)$ are, respectively, the momentum integration of the dissociation probability $W_{g\Psi}^{c\bar{c}}$ multiplied by gluon thermal distribution f_g and the regeneration probability $W_{c\bar{c}}^{g\Psi}$ multiplied by charm quark distribution f_c [15]. $W_{g\Psi}^{c\bar{c}}$ is calculated with perturbative Coulomb potential [24] and $W_{c\bar{c}}^{g\Psi}$ can be obtained from $W_{g\Psi}^{c\bar{c}}$ via detailed balance. Since the experimentally measured D-meson flow is comparable with light hadron flow [25], we assume charm quark thermalization and choose the density $f_c(q, x) = T_A(\mathbf{x}_t)T_B(\mathbf{x}_t - \mathbf{b}) \cosh \eta / \tau d\sigma_{NN}^{c\bar{c}} / d\eta f_q(q, x)$, where f_q is the normalized charm quark thermal distribution, $d\sigma_{NN}^{c\bar{c}} / d\eta$ the rapidity distribution of charm quark pairs produced in NN collisions, $\mathbf{b}$ the impact parameter, and T_A and T_B are the thickness functions of the two colliding nuclei defined as $T(\mathbf{x}_t) = \lim_{z_1 \rightarrow -\infty, z_2 \rightarrow \infty} T(\mathbf{x}_t, z_1, z_2)$ with

$T(\mathbf{x}_t, z_1, z_2) = \int_{z_1}^{z_2} dz \rho(\mathbf{x}_t, z)$ and the Woods-Saxon nuclear density profile $\rho(\mathbf{x})$.

Since the hadronic phase appears later in the evolution of the fireball when the density of the system is lower compared to the early hot and dense period, we have neglected the charmonium production and suppression in hadron gas. On the other hand, from the lattice QCD simulation [26] on charmonium spectral function at finite temperature, J/ψ collapses at a dissociation temperature $T_d^{J/\psi}$, which corresponds to the idea of sequential suppression [27]. Taking into account these two aspects, the dissociation and regeneration processes happen only in the temperature region $T_c < T < T_d^\Psi$ of QGP, where T_c is the critical temperature of deconfinement phase transition. Outside this region there are $\alpha_\Psi = \beta_\Psi = 0$ for $T < T_c$ and $\alpha_\Psi = \beta_\Psi = \infty$ for $T > T_d^\Psi$.

The transport equation has been solved analytically and the result is shown as

$$f_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{x}_t, \eta, \tau) = f_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{X}_\Psi(\tau_0), H_\Psi(\tau_0), \tau_0) e^{-\int_{\tau_0}^{\tau} d\tau' \alpha_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{X}_\Psi(\tau'), H_\Psi(\tau'), \tau') / \Delta(\tau')} \\ + \int_{\tau_0}^{\tau} d\tau' \beta_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{X}_\Psi(\tau'), H_\Psi(\tau'), \tau') / \Delta(\tau') e^{-\int_{\tau'}^{\tau} d\tau'' \alpha_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{X}_\Psi(\tau''), H_\Psi(\tau''), \tau'') / \Delta(\tau'')} \quad (2)$$

with

$$\mathbf{X}_\Psi(\tau') = \mathbf{x}_t - \mathbf{v}_t^\Psi [\tau \cosh(y_\Psi - \eta) - \tau' \Delta(\tau')], \\ H_\Psi(\tau') = y_\Psi - \text{arcsinh}(\tau / \tau' \sinh(y_\Psi - \eta)), \\ \Delta(\tau') = \sqrt{1 + (\tau / \tau')^2 \sinh^2(y_\Psi - \eta)}. \quad (3)$$

The first and second terms on the right-hand side of the solution (2) indicate the contributions from the initial production and continuous regeneration, respectively, and both suffer anomalous suppression in the medium. Since the collision time for NN interactions at RHIC energy is about 0.1 fm/c and less than the starting time τ_0 of the medium evolution which is about 0.5 fm/c, the nuclear absorption and Cronin effect [28] for the initially produced charmonia have ceased before the QGP evolution and can be reflected in the initial distribution of the transport equation (1). From the PHENIX d+Au data, the cold nuclear matter effect is smaller at RHIC than those observed at lower energies [29]. Considering further a finite formation time of charmonia which is about 0.5 fm/c and larger than the collision time, the nuclear absorption can approximately be neglected for the calculations at high energies [16]. In this case, the initial distribution $f_\Psi(\mathbf{p}_t, y_\Psi, \mathbf{x}_t, \eta, \tau_0)$ is calculated according to the Glauber model with the J/ψ production cross section $d\sigma_{NN}^\Psi / (p_t dp_t dy_\Psi)$ in NN collisions and the modification of the Cronin effect [28]. The coordinate shifts $\mathbf{x}_t \rightarrow \mathbf{X}_\Psi$ and $\eta \rightarrow H_\Psi$ in the solution (2) reflect the leakage effect in the transverse and longitudinal directions.

The local temperature, baryon chemical potential and fluid velocity are determined by hydrodynamic equa-

tions and control the suppression and regeneration region through the gluon and quark thermal distributions. Similar to the three dimensional transport equation for Ψ motion, a self-consistent treatment for the calculation of J/ψ rapidity dependence needs three dimensional hydrodynamic equations. Considering the fact that the experimentally observed forward rapidity $1.2 < y < 2.2$ is still located at the central plateau of the Bjorken hydrodynamics at RHIC energy [23], we can, for simplification, neglect the rapidity dependence of the fluid and take the transverse hydrodynamic equations [15, 16] at $y = 0$ for the QGP evolution at both mid and forward rapidity. To close the equations, we take the equation of state [30] of ideal gases of partons and hadrons with a first order phase transition at T_c , and the initial condition for the hydrodynamics at RHIC is the same as in Ref [15, 16].

We now fix the charmonium and charm quark distributions in NN collisions. With the precise transverse momentum and rapidity distributions for J/ψ measured in proton-proton collisions at RHIC energy [31], the distribution $d\sigma_{NN}^\Psi / (p_t dp_t dy_\Psi)$ can be parameterized as

$$\frac{d\sigma_{NN}^\Psi}{p_t dp_t dy_\Psi} = \frac{2(n-1)}{D(y_\Psi)} \left(1 + \frac{p_t^2}{D(y_\Psi)}\right)^{-n} \frac{d\sigma_{NN}^\Psi}{dy_\Psi} \quad (4)$$

with $n = 6$, $D(y_\Psi) = \langle p_t^2 \rangle_{NN} (n-2) (1 - y_\Psi^2 / Y_\Psi^2)$, the maximum rapidity $Y_\Psi = \text{arccosh}(\sqrt{s_{NN}} / (2m_\Psi))$, the averaged transverse momentum square $\langle p_t^2 \rangle_{NN} = 4.14 \text{ (GeV/c)}^2$ and a double Gaussian distribution $d\sigma_{NN}^\Psi / dy_\Psi$ [31]. At LHC energy, from the CDF experiment and CEM calculation [32, 33], we take $n = 4$,

$\langle p_t^2 \rangle_{NN} = 12 \text{ (GeV/c)}^2$ and $d\sigma_{NN}^{\Psi}/dy_{\Psi} = 2 \mu b$. Taking into account the Cronin effect [28], $\langle p_t^2 \rangle_{NN}$ is replaced by $\langle p_t^2 \rangle_N = \langle p_t^2 \rangle_{NN} + a(T_A(\mathbf{x}_t, -\infty, z_A) + T_B(\mathbf{x}_t - \mathbf{b}, z_B, \infty))$. The second term is from the multiply scattering of the two gluons with nucleons before they fuse into a Ψ , and the constant a is adjusted to the data for pA collisions [15, 34].

For $d\sigma_{NN}^{c\bar{c}}/d\eta$ at RHIC, there is a large experimental uncertainty and the difference among theoretical model estimations is also significant [35]. We take a Gaussian distribution $d\sigma_{NN}^{c\bar{c}}/d\eta = d\sigma_{NN}^{c\bar{c}}/d\eta|_{\eta=0} e^{-\eta^2/\eta_0^2}$ with $d\sigma_{NN}^{c\bar{c}}/d\eta|_{\eta=0} = 120 \mu b$ which agrees with the experimental data [35] and a parameter η_0 determined by the assumption $(d\sigma_{NN}^{c\bar{c}}/d\eta|_{\eta=1.7})/(d\sigma_{NN}^{c\bar{c}}/d\eta|_{\eta=0}) = 1/3$ which is in between the maximum and minimum model calculations [35]. At LHC, we take $d\sigma_{NN}^{c\bar{c}}/d\eta$ directly from the PYTHIA simulation [36].

Taking the dissociation temperatures $T_d^{J/\psi} = 1.9 T_c$, $T_d^{\chi_c} = T_d^{\psi'} = T_c = 165 \text{ MeV}$ and the J/ψ fractions from direct production and χ_c and ψ' decay as 6 : 3 : 1, we integrated the J/ψ distribution $f_{J/\psi}$ on the hadronization hyper surface and calculated the J/ψ R_{AA} and $\langle p_t^2 \rangle$ at mid and forward rapidity in heavy ion collisions at RHIC and LHC energies. In the upper panel of Fig.1 we show the ratio of forward to mid rapidity R_{AA} as a function of N_p at RHIC. The new experimental finding of $R_{AA}^{\text{forward}}/R_{AA}^{\text{mid}} < 1$ is difficult to be understood in models with only initial production mechanism, since the suppression in forward region should be smaller than or at most the same as that in central region. In our calculation with the assumption of the same medium for the two rapidity regions, the ratio with only initial production is almost a constant, the small deviation is from the rapidity dependence of the J/ψ transverse momentum distribution in NN collisions, see $D(y_{\psi})$ in (4). For the case with only regeneration, the ratio becomes a strict constant and is determined by the cross sections σ_{Ψ} and $\sigma_{c\bar{c}}$ in NN collisions. Since the regeneration contribution at forward rapidity is smaller than that at midrapidity, the total ratio is less than the value with only initial production in the whole N_p region. However, it can not reach the regeneration limit, because at RHIC the maximum fraction of regeneration is only around 50% [15].

The ratio of forward to mid rapidity $\langle p_t^2 \rangle$ is shown in the lower panel of Fig.1. Again it is a constant in the case with only initial production or regeneration. While the $\langle p_t^2 \rangle$ is dominated by the initial production at forward rapidity, both initial production and regeneration affect the midrapidity region. In addition, the initially produced J/ψ s are harder in momentum distribution. Therefore, the total ratio is always larger than the limiting value set by the initial production. It starts with the value of $\langle p_t^2 \rangle_{NN}^{\text{forward}}/\langle p_t^2 \rangle_{NN}^{\text{mid}}$ and goes up with increasing N_p monotonously.

While the J/ψ production at RHIC is governed by the competition between the initial production and regeneration, the situation at LHC is very different. At LHC, the




FIG. 1: The ratio of forward to mid rapidity R_{AA} and the ratio of forward to mid rapidity $\langle p_t^2 \rangle$ at RHIC. The lines are theoretical calculations and the data are from [17, 19].




FIG. 2: The ratio of forward to mid rapidity R_{AA} , the ratio of forward to mid rapidity $\langle p_t^2 \rangle$ and the ratio of forward to mid rapidity S_{AA} in Pb+Pb collisions at LHC.

fireball is hotter, larger, and longer lived, almost all the initially produced J/ψ s are eaten up by the QGP. On the other hand, there are about ten times more charm quarks at LHC than that at RHIC. As a result, the regeneration becomes much more important and dominates the J/ψ production in heavy ion collisions at LHC. We show our prediction for the two ratios in Fig.2. The midrapidity and forward rapidity regions are defined as $|y| < 0.9$ and $y \in [2.5, 4]$. With increasing N_p , both ratios start from the initial production limit and reaches the regeneration limit fast. At large N_p , the two ratios are almost fully controlled by the regeneration. The rapid change of the two ratios at low N_p is due to the important contribution from the initial production in peripheral collisions, like the case in central collisions at RHIC.

When the J/ψ regeneration becomes dominant, we can use the quantity $S_{AA} = N_{J/\psi}/N_D^2$ to describe the medium effect, where $N_{J/\psi}$ and N_D are, respectively, the J/ψ number and D-meson number per unit rapidity. Since the number of initially produced J/ψ s is proportional to the number of binary collisions N_c and the number of regenerated J/ψ s to N_c^2 , we have $S_{AA} \sim 1/N_c$ in the initial production dominant region and $S_{AA} \sim 1$

in the regeneration dominant region. The ratio of forward to mid rapidity S_{AA} at LHC is shown in Fig.2. In contrast to the ratio of R_{AA} which reaches unity in the limit of small N_p , the ratio of S_{AA} approaches to unity in the limit of large N_p .

In summary, we have constructed a three dimensional transport model to describe the rapidity dependence of J/ψ production in relativistic heavy ion collisions. To understand the production and suppression mechanisms, we need to investigate simultaneously the J/ψ yield and transverse momentum distribution. At RHIC, the competition between the initial production and regeneration explains well the rapidity dependence of J/ψ R_{AA} and $\langle p_t^2 \rangle$, namely $R_{AA}^{\text{forward}}/R_{AA}^{\text{mid}} < 1$ at any centrality and $\langle p_t^2 \rangle^{\text{forward}}/\langle p_t^2 \rangle^{\text{mid}} > 1$ for semi-central and central collisions. At LHC, the two ratios approach to the corresponding regeneration limits already in semi-central collisions, due to the fast dissociation of the initially produced charmonia and the dominance of the regeneration.

Acknowledgement: The work is supported by the NSFC Grant 10735040, the 973-project 2006CB921404 and 2007CB815000, and the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

-
- [1] T.Matsui and H.Satz, Phys. Lett. **B178**, 416(1986).
 - [2] C.Gerschel and J.Hufner, Ann. Rev. Nucl. Part. Sci. **49**, 255(1999).
 - [3] R.Vogt, Phys. Rept. **310**, 197(1999).
 - [4] M.C.Abreu *et al.*, [NA50 Collaboration], Nucl. Phys. **A610**, 404c(1996).
 - [5] J.-P.Blaizot and J.-Y.Ollitrault, Phys. Rev. Lett. **77**, 1703(1996).
 - [6] A.Polleri *et al.*, Phys. Rev. **C70** 044906(2004).
 - [7] A.Capella *et al.*, Phys. Lett. **B393**, 431(1997).
 - [8] E.Bratkovskaya *et al.*, Phys. Rev. **C71** 044901(2005).
 - [9] J.Hüfner and P.Zhuang, Phys. Lett. **B559**, 193(2003).
 - [10] P.Braun-Munzinger and J. Stachel, Phys. Lett. **B490**, 196(2000).
 - [11] M.I.Gorenstein *et al.*, Phys. Lett. **B509**, 277(2001).
 - [12] L.Grandchamp and R.Rapp, Phys. Lett. **B523**, 60(2001).
 - [13] V.Greco, C.M.Ko and R.Rapp, Phys. Lett. **B595**, 202(2004).
 - [14] R.L.Thews and M.L.Mangano, Phys. Rev. **C73**, 014904(2006).
 - [15] L.Yan, P.Zhuang and N.Xu, Phys. Rev. Lett. **97**, 232301(2006).
 - [16] Y.Liu *et al.*, J. Phys. **G36**, 064057(2009).
 - [17] A.Adare *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **98**, 232301(2007).
 - [18] B.I.Abelev, *et al.*, [STAR Collaboration], arXiv:0904.0439.
 - [19] J.Lajoie, [PHENIX Collaboration], J. Phys. **G34**, S191(2007).
 - [20] D.Kharzeev *et al.*, arXiv:0809.2933.
 - [21] A.Andronic *et al.*, J.Phys. **G35**, 104155(2008).
 - [22] X.Zhao and R.Rapp, arXiv:0810.4566.
 - [23] I.G.Bearden *et al.*, [BRAHMS Collaboration], Phys. Rev. Lett. **88**, 202301(2002).
 - [24] G. Bhanot, M.E. Peskin, Nucl. Phys. **B156**, 391(1979).
 - [25] A.Adare *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **98**, 172301(2007).
 - [26] M.Asakawa and T.Hatsuda, Nucl. Phys. Proc. Suppl. **129**, 584(2004).
 - [27] H.Satz, Nucl. Phys. **A783**, 249(2007).
 - [28] S.Gavin and M.Gyulassy, Phys. Lett. **B214**, 24 1(1988); J.Hüfner, Y.Kurihara and H.J.Pirner, Phys. Lett. **B215**, 218(1988).
 - [29] S.S.Adler *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **96**, 012304(2007).
 - [30] J.Sollfrank *et al.*, Phys. Rev. **C55**, 392(1997).
 - [31] A.Adare *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **98**, 232002(2007).
 - [32] B.Alessandro *et al.*, [ALICE Collaboration], J. Phys. **G32**, 1295(2006).
 - [33] A.Accardi *et al.*, hep-ph/0308248.
 - [34] X.Zhao and R.Rapp, Phys. Lett. **B664**, 253(2008).
 - [35] Y.Zhang, J. Phys. **G35**, 104022(2008).
 - [36] N.Carrer and A.Dainese, hep-ph/0311225.